



TITLE:

Enzymic Dehydrochlorination of Trichlorofon by the Digestive Juice of the Silkworm, Bombyx mori L

AUTHOR(S):

SUGIYAMA, Hiroshi; SHIGEMATSU, Hajime

CITATION:

SUGIYAMA, Hiroshi ...[et al]. Enzymic Dehydrochlorination of Trichlorofon by the Digestive Juice of the Silkworm, Bombyx mori L. 防虫科学 1969, 34(2): 79-85

ISSUE DATE:

1969-05-31

URL:

<http://hdl.handle.net/2433/158573>

RIGHT:

Enzymic Dehydrochlorination of Trichlorofon by the Digestive Juice of the Silkworm, *Bombyx mori* L. Hiroshi SUGIYAMA and Hajime SHIGEMATSU (The Sericultural Experiment Station, Suginami-ku, Tokyo). Received March 28, 1969. *Botyu-Kagaku* 34, 79, 1969.

10. 家蚕幼虫消化液によるトリクロロフォンの酵素的脱塩酸作用. 杉山 浩・重松 孟 (農林省蚕糸試験場) 44. 3. 28 受理.

家蚕幼虫の消化液によるトリクロロフォンの DDVP への非酵素的あるいは酵素的移行を紫外分光、薄層クロマトグラフィ、赤外分光および若干の化学反応による方法で調べた。トリクロロフォンは消化液によらない弱アルカリ溶液中で非酵素的にすみやかに DDVP に変わる。中性、弱酸性ではその移行はいちじるしく少ない。しかし、消化液中でトリクロロフォンを作用すると、反応液の pH が中性ないし弱酸性であっても急速に DDVP に移行した。この移行はおもに酵素的な反応で、その反応の至適 pH は 7 近辺にあった。食桑中の幼虫の消化液の pH—9 付近では、トリクロロフォンから DDVP への移行は非酵素反応と酵素反応の両者が関与すると考えられた。DEAE-セルロースカラムでこの酵素を分画することが可能で、消化液たんばくから分離して得られた。濾紙電気泳動においてもこの活性たんばくは移動度がもっとも速く消化液たんばくから分離した。

トリクロロフォンの分解産物として DDVP のほかにその近縁化合物が生成していた。

The toxicity of trichlorofon (*O, O*-dimethyl 2, 2, 2-trichloro 1-hydroxyethylphosphonate) is higher in the oral application than in the topical application to the silkworm larva²⁾. In the preceding paper it was recovered that trichlorofon was rapidly converted into DDVP (*O, O*-dimethyl 2, 2-dichlorovinylphosphate), which is a powerful inhibitor of cholinesterase, in the digestive juice through dehydrochlorination, but not in the hemolymph of the silkworm larva *in vitro*³⁾. It has been suggested that the insecticide was rapidly dehydrochlorinated in a weak alkaline solution^{3, 4)}. We found the occurrence of active conversion of trichlorofon even with use of the neutral digestive juice. Therefore, we will pay our attentions to a problem if the dehydrochlorination of trichlorofon is dependent on an enzymic reaction or simply on the alkalinity of the digestive juice.

Materials and Methods

The digestive juice from the larvae of the silkworm, *Bombyx mori*, of a hybrid between the strains Nichi 124 and Shi 124 was collected on the fourth to sixth days of the fifth instar, after one day of fasting, with vomiting by an electric stimulation. The juice was stored at -20°C until use. After thawing, the supernatant fluid resulted after the centrifugation of the juice for 10 minutes at 1,000 G was subjected to the assay.

Trichlorofon was purified from 80% soluble

powdery formulation by recrystallization from *n*-hexane. A technical DDVP of 98% purity was dissolved in ether and washed with distilled water for several times. When the solvent was removed under a diminished pressure, DDVP was obtained as a viscous liquid and stored in an ampule until use.

Experiments on the non-enzymic conversion of trichlorofon into DDVP were carried out with a mixture of 0.8 ml of a definite pH in 0.5M Tris (hydroxyethylaminomethane)-NaOH-HCl buffer solution and 2.57 mg trichlorofon in 0.2 ml of distilled water at 28°C. The reaction products were once extracted from the reaction mixture with 4 ml of acetonitrile. Before the determination with spectrophotometer, acetonitrile extracts were cleaned by the centrifugation.

Experiments with the digestive juice of larva were carried out by mixing 0.2 ml of the juice, 0.5 ml of Tris-HCl buffer, 0.1 ml of 10⁻²M KCN solution and 2.57 mg trichlorofon in 0.2 ml and incubated at 35°C. The reaction products were extracted with acetonitrile as described above. For an anaerobic reaction, a Thunberg tube was used after sucking air for 3 minutes by an aspirator.

Acetonitrile extracts were optically examined with ultraviolet absorptions. Careful attention should be paid to shifting of absorption maximum with different kind of solvents or amounts of the digestive juice added. So far we examined, DDVP

in acetonitrile solution showed an absorption maximum at $212\sim 214\text{m}\mu$, while in acetonitrile including Tris-NaOH-HCl buffer, the absorption maximum of DDVP shifted to $218\text{m}\mu$. Moreover, in the solution added with the digestive juice, the absorption maximum shifted to $220\sim 230\text{m}\mu$. Silica gel thin-layer chromatography, developing by benzene-methanol (9:1, v/v), was employed to separate decomposition products of trichlorofon. Visible spots of chromatogram under ultraviolet light were extracted with acetonitrile and assayed for ultraviolet and infrared absorptions. By this solvent system, Rf values for trichlorofon, DDVP, and another decomposition product were $0.0\sim 0.10$, $0.55\sim 0.80$, and $0.20\sim 0.35$, respectively.

Results

Determination of DDVP by Ultraviolet Absorption: The C=C double bond of DDVP is responsible to the absorption of ultraviolet light. Possibility of the determination of DDVP was examined by the ultraviolet absorption in acetonitrile. Various amount of DDVP dissolved in 0.5 ml of acetonitrile and 1 ml of 0.5 M Tris-HCl buffer (pH 7.0) was vigorously shaken with 4 ml of acetonitrile and then centrifuged for 5 minutes at 1,000 G. Acetonitrile layer was assayed for optical absorption at $218\text{m}\mu$. The optical absorption of DDVP was linearly related to the concentration of the compound (A curve in Fig. 1). Recovery of DDVP from the digestive juice was tested as follows: various amounts of DDVP dissolved in 0.5 ml of acetonitrile were mixed with 0.2 ml of the juice and 0.8 ml of the buffer (pH 7.0), and DDVP was extracted with 4 ml of acetonitrile by vigorous shaking as mentioned above. Since DDVP in the acetonitrile extract showed absorption maximum at $230\text{m}\mu$, the optical density at $230\text{m}\mu$ was plotted against the concentration of DDVP initially added. Results were shown with curve B in Fig. 1. Recovery of DDVP was $64\sim 67\%$ in this case. Shift of the absorption maximum and change of recovery of DDVP had to be pre-examined in experiments with use of the digestive juice. As previously mentioned⁹⁾, trichlorofon was converted into at least two compounds, detectable on the thin-layer chromatogram, which seem to be DDVP and its analogue, and the latter compound showed also

ultraviolet absorption as described later. Therefore we had to temporarily express the amount of conversion products of trichlorofon as DDVP equivalence calculated from the lines of Fig. 1. Trichlorofon in acetonitrile did not absorb the ultraviolet light at the range between 200 and $260\text{m}\mu$ as reported by Tokumitsu⁶⁾.




Fig. 1. Spectrophotometric determination of DDVP. A: in the absence of the digestive juice, measured at $218\text{m}\mu$, B: in the presence of the digestive juice, measured at $230\text{m}\mu$.

Non-enzymic Conversion of Trichlorofon into DDVP: Trichlorofon was rapidly converted into DDVP in an alkaline solution. The time course of DDVP formation from trichlorofon under various pH was shown in Fig. 2. The higher in pH value of the solution, the higher in the initial velocity of the conversion of trichlorofon. However, trichlorofon in an alkaline solution was not completely degraded. For instance, DDVP produced at pH 9.8 in 60 minutes-reaction amounted to 1.05 mg, only about half of trichlorofon initially added, and then slightly fell down in accordance to reaction time; it was not clear whether DDVP was further converted into other products or not. At pH 9.0 it took almost 100 minutes to arrive at the same maximum level of DDVP production at pH 9.8. In the reactions at pH 8.1 and 7.1, the reaction velocity decreased remarkably.

Conversion of Trichlorofon into DDVP in the Digestive Juice in Vitro: When the digestive juice was left to stand after centrifugation, pH was reduced from about 10 to 7 within 30 minutes. Since it was difficult to keep constant the pH of




Fig. 2. The amount of DDVP converted non-enzymically from 2.57 mg of trichlorofon in various pH solutions.

the digestive juice with a dilute buffer, 0.5 M Tris-HCl buffer was employed in this investigation.

As shown in Table 1, activity was not significantly detected in the reaction under aerobic condition, but clearly observed under anaerobic condition or by an addition of KCN. Optimum concentration of KCN was found to be around 1×10^{-3} M (Table 2). Reduced glutathion or NADPH₂ did not accelerate the activity. Under the aerobic condition of the digestive juice, the conversion

Table 1. Activity of the digestive juice on dehydrochlorination of Trichlorofon under aerobic and anaerobic conditions at pH 7.0. The reaction mixture was prepared according to the methods and incubation was allowed to proceed for 30 minutes at 35°C.

Reaction condition	Amount of DDVP formed (µg)
aerobic	60
anaerobic	130
aerobic+KCN(1×10^{-3} M)	225




Fig. 3. Time course of DDVP formation from trichlorofon in the digestive juice at pH 7.0. Value of DDVP formation at each reaction time was subtracted by that obtained at zero time.

Table 2. Effect of KCN addition to the digestive juice on an aerobic reaction at pH 7.0. Reaction condition was the same as Table 1.

Reaction condition	Amount of DDVP formed (µg)
KCN added, 1×10^{-2} M	195
KCN added, 1×10^{-3} M	475
KCN added, 1×10^{-4} M	40
KCN not added	trace
Digestive juice omitted (no KCN)	15
With boiled digestive juice (no KCN)	trace




Fig. 4. Relationship between the amount of DDVP formed from trichlorofon and the amount of the digestive juice added during 5 minutes reaction at pH 7.0. Value of DDVP formation with each amount of the digestive juice was subtracted by that of the zero-time reaction.




Fig. 5. The shift of ultraviolet absorption spectra in accordance with the amount of the digestive juice after the 5 minutes reaction at pH 7.0.

seemed to be inhibited somehow. When the digestive juice was boiled for 10 minutes in a heating bath, trichlorofon was not converted.

The reaction in the digestive juice proceeded linearly for at least 30 minutes (Fig. 3) and the activity seemed to be proportional to the amount of the digestive juice within 5 minutes (Fig. 4).

Absorption maximum of the reaction product

was shifted to longer wave length with use of larger volume of the digestive juice (Fig. 5).

The conversion of trichlorofon into DDVP in the digestive juice was examined under various pH values ranging from 5 to 11 (Table 3). Tris-HCl buffer was employed, but over pH 11 and below 7, NaOH and HCl were added to give appropriate values, respectively. The reaction without the digestive juice was set as a reference. It was noticed at first from the results in Table 3 that the values of optical absorption at zero time reactions were not ignorant, depending upon pH values, and were supposed to be formed during the period of extraction of the reaction products. These values responded to the non-enzymic reaction shown in Fig. 2.

A remarkable production of DDVP in these experiments was observed at pH 7 to 8. Amounts of DDVP formed by non-enzymic conversion of trichlorofon were observed to be 230 and 65 µg at pH 8.3 and 7.0, respectively, while in enzymic reaction 555 and 425 µg of DDVP were formed at pH 8.2 and 6.3, respectively. Larger amount of produced DDVP in enzymic reaction at a high pH range, for example, 780 µg at pH 9.0, might be resulted by an additional effect of non-enzymic and enzymic conversions. However, the enzymic action related to the dehydrochlorination of trichlorofon in the digestive juice was not effective above pH 10. This enzyme seemed to have the optimum pH at neutral or rather slightly acidic.

Attempts for Purification of the Enzyme: Preliminary trials for purification of the enzyme were carried out. This enzyme could be fractionated by columns of either Sephadex G-50 or G-100. Using the column of DEAE-cellulose equilibrated by 0.05M Tris buffer (pH 7.4), fractionated by NaCl solution with the linear gradient of its concentration from 0 to 1.0M, the active fractions were eluted with around 0.6M salt concentration, after major peaks of juice protein, as presented in Fig. 6.

Zymography was attempted to visualize the active band of protein on paper strip through electrophoresis. A small amount of the digestive juice or the active eluate from the column was placed on strips of Toyo filter paper no. 51A and the electrophoresis was performed for 3 hours

Table 3. Conversion of trichlorofon into DDVP in the digestive juice at various pH.

The reaction mixture was prepared according to the methods. Incubation the was allowed to proceed for 15 minutes at 35°C.

pH of the reaction mixture	Optical density at zero time	Increase of optical density at 15 min.-reaction	Amount of DDVP formed in 15 min.
(A) Reactions without the digestive juice			
11.4	1.09 (223m μ) ^a	0.420(227m μ) ^a	335 μ g
10.0	0.77 (222m μ)	0.620(225m μ)	495
9.0	0.396(221m μ)	0.614(223m μ)	490
8.3	0.306(219m μ)	0.292(220m μ)	232
7.5	0.243(218m μ)	0.095(217m μ)	75
7.0	0.218(218m μ)	0.087(217m μ)	65.5
(B) Reactions with the digestive juice			
11.2	1.08 (228m μ)	0.223(234m μ)	280 μ g
9.9	1.08 (230m μ)	0.233(235m μ)	290
9.0	0.522(226m μ)	0.653(230m μ)	780
8.2	0.288(224m μ)	0.462(227m μ)	555
7.2	0.186(222m μ)	0.349(223m μ)	425
6.3	0.223(221m μ)	0.350(223m μ)	425
5.4	0.136(223m μ)	0.308(221m μ)	375
(C) Rough estimation of DDVP formation owed to enzymic action of the digestive juice (B-A)			
11.3			— ^b μ g
10.0			— ^b
9.0			290
8.3			323
7.1			360
6.3			370 ^c
5.4			340 ^c

^a Wave-length of absorption maximum was shown in parentheses

^b Enzymic conversion seems not to be accounted

^c Amount of non-enzymic conversion was extrapolated from data of (A)

with 1.0mA/cm in the cold room, using Veronal buffer (μ =0.1, pH 8.6). After the run, the paper strip was gently sprayed by 0.1M Tris buffer, pH 6.6, and coated by trichlorofon solution, followed by incubation for 30 minutes at 35°C. And then, the paper was dried and colored by the adequate methods for DDVP or protein. The results suggested that the active band had the highest mobility, aparted from major bands of juice protein.

Another Derivative of DDVP: Another derivative of DDVP suggested in previous experiments⁵⁾ was separated by thin-layer chromatography on silica gel plate of 500 μ thickness. The substance showed R_f value 0.20~0.35 with the benzene-

methanol solvent system and was extracted from the plates by acetonitrile, and was assayed by ultraviolet and infrared spectrophotometry and chemical reaction. As shown in Fig. 7, a large absorption at 212m μ which was characteristic for DDVP was recognized in ultraviolet absorption spectrum of this compound. Infrared spectra of DDVP and of this compound were shown in Fig. 8. As compared with the spectrum of DDVP, this compound showed the following natures: OH vibration occurred near 3,200cm⁻¹; two new bands absorption at 1,122 and 1,075cm⁻¹ appeared instead of 1,150 and 1,050cm⁻¹; a characteristic absorption of DDVP at 980cm⁻¹ was not found, but new absorption bands were found at 967 and




Fig. 6. Elution pattern from DEAE-cellulose column of an enzyme catalyzing dehydrochlorination of trichlorofon from the digestive juice. A linear NaCl gradient (0 to 1.0M) was employed and fractions of 3ml were collected. —: protein, ...: enzyme activity, ---: NaCl concentration.




Fig. 7. Ultraviolet absorption spectra of DDVP and of another derivative separated by thin-layer chromatography. I: extract from Rf 0.55~0.80 region (DDVP), II: extract from Rf 0.20~0.35 region; chromatoplate, silica gel, 500 μ in thickness; developing solvent system, benzene-methanol (9:1, v/v).

943 cm^{-1} suggesting the presence of the vinyl moiety; in the 900-700 cm^{-1} range a band at 855 cm^{-1} seen in the spectrum of DDVP was absent from that of the derivative, whilst a new band appeared at 745 cm^{-1} ; and a strong absorption at 1,280 cm^{-1} associated with P=O vibration was found in the spectrum of both DDVP and the




Fig. 8. Infrared absorption spectra of DDVP and of another derivative separated by thin-layer chromatography. I: technical DDVP 98%, II: extract from Rf 0.55~0.80 region, III: extract from Rf 0.20~0.35 region, KBr disc method was applied.

derivative. The presence of a non-conjugated C=C bond was further supported by the strong absorption at 1,650 cm^{-1} in infrared spectra both for DDVP and the another decomposed material, and by the chemical reaction with potassium permanganate in sodium carbonate where decoloration occurred by the derivative as well as by DDVP. These data suggested that both the

derivative and DDVP possessed C=C bond in the molecules. From these indications, it seems likely to assume that this compound is not DDVP, but some closed compound to DDVP.

Discussion

The pH value of the digestive juice of silkworm larva is about 9 in feeding period and much higher in fasting condition¹⁾, while that of the hemolymph of the larva is around 6.5. It is, therefore, reasonable to say that the conversion of trichlorofon is more rapid in the digestive juice than in the hemolymph owing to the difference of pH, as explained in the previous paper²⁾.

In the present experiments, it was shown that trichlorofon converted into DDVP even near pH 7 in the digestive juice, while the formation of DDVP was not significant in the non-enzymic reaction at this pH. This result indicates a possibility that an enzymic action takes a part in the dehydrochlorination of trichlorofon.

The conversion of trichlorofon into DDVP in the digestive juice seemed to be carried mainly by a non-enzymic reaction in high alkaline conditions, but by an enzymic reaction in low alkaline or neutral conditions. Miyamoto⁴⁾ pointed that trichlorofon was converted into DDVP by non-enzymic transformation in neutral and acidic conditions and the resulted DDVP worked as an inhibitor of acetylcholinesterase in the fly head *in vitro*. Under neutral or acidic condition, from our experimental results, it is, however, doubtful to participate significantly the non-enzymic reaction to the conversion of trichlorofon. Certainly, in the digestive juice of the silkworm larva, non-enzymic transformation may not be ignored, even if there should be the enzymic reaction. At any rate, in the digestive juice, trichlorofon applied orally can be transformed into DDVP under wide range of pH by both non-enzymic and enzymic actions. Studies on the enzyme which is responsible for the dehydrochlorination of trichlorofon in the digestive juice is now under way.

Summary

The non-enzymic or enzymic conversion of

trichlorofon into DDVP in the digestive juice of the silkworm larva was studied by means of ultraviolet spectrophotometry, thin-layer chromatography, infrared spectrophotometry and some chemical reactions. Trichlorofon was rapidly converted into DDVP in a mild alkaline solution, non-enzymically, but very slowly in neutral or slightly acidic condition. On the other hand, in the digestive juice of the silkworm larva trichlorofon rapidly changed to DDVP under a broader pH range of conditions, even in neutral or slightly acidic. It was shown that around neutral conditions, the conversion in the digestive juice progressed mainly enzymically. The enzyme activity concerned this conversion was recognized in the biological fluid, the pH optimum being around 7 or slightly lower. Near pH 9, which is normal pH of the digestive juice of silkworm larva feeding mulberry leaves, both non-enzymic and enzymic reactions seemed to participate in the conversion of trichlorofon. Enzymic activity was negligible above pH 10. Using the column of DEAE-cellulose, the active protein fractions were obtained. The active protein could be separated from juice proteins on zymograms of paper electrophoresis. In addition to DDVP, another decomposed compound of trichlorofon was formed during the reaction and this compound was partly characterized.

Acknowledgement: The authors are deeply indebted to Prof. Z. Kuwana of Bunka Women's College for his encouragement and valuable advices in this work.

Literature Cited

- 1) Fujii, O.: *Biochemical Studies on the Silkworm Flacherry*. pp.36, Sangyo-shimpo-sha, Tokyo (1936).
- 2) Kuwana, Z., Nakamura, S. and Sugiyama, H.: *Bull. Sericul. Exp. Sta.*, 22, 123 (1967).
- 3) Metcalf, R. L., Fukuto, T. R. and March, R. B.: *J. Econ. Entomol.*, 59, 44 (1959).
- 4) Miyamoto, J.: *Botyu-Kagaku*, 24, 130 (1959).
- 5) Sugiyama, H.: *Appl. Ent. Zool.*, 1, 197 (1966).
- 6) Tokumitsu, T.: *Kogyo-Kagaku Zasshi*, 67, 649 (1964).